

English

## 

(43) International Publication Date 18 January 2007 (18.01,2007)

## (51) International Patent Classification: C07K 16/28 (2006.01) A61K 39/395 (2006.01)

- A61K 39/39 (2006.01) A61P 35/00 (2006.01)
- (21) International Application Number: PCT/US2006/025711
- (22) International Filing Date: 30 June 2006 (30.06.2006)
- (25) Filing Language:
- (26) Publication Language: English
- (30) Priority Data: 60/697,082
- 7 July 2005 (07.07.2005)
- (71) Applicants (for all designated States except US): COLEY PHARMACEUTICAL GROUP, INC. [US/US]: 93 Worcester Street, Suite 101, Wellesley, MA 02481 (US). PFIZER INC. [US/US]: 235 East 42nd Street, New York, NY 10017 (US).
- (72) Inventors; and
- (1/2) Inventors; and (1/2) Inventors/Applicants (for US only): READETT, David, Robert, John (IBI/US); 10 New London Road, Mystic, CT 03535 (US), JUNGNELLUS, Jarf, UIE, Birger [SE/US]; 178 Long Wharf Drive, Mystic, CT 0355 (US), GOMEZ-NOVARRO, Jesus [ES/US]; 18 Whitehall Landing, Mystic, CT 03555 (US), HANSON, Dougles, C. [US/US]; 3 Acom Drive, Niantic, CT 03537 (US), KRIEGA, Arthur, M. [US/US]; 173 Winding River Road, Wellesley, MA 02482 (US)

- (10) International Publication Number WO 2007/008463 A2
- (74) Agent: TREVISAN, Maria, A.; Wolf, Greenfield & Sacks, P.C., 600 Atlantic Avenue, Boston, MA 02210 (US).
- (81) Designated States (unless otherwise indicated for every kind of national protection available); R.E., AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, BG, GS, FI, GB, GD, GE, GH, GM, HN, HR, HU, DI, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LC, LR, LIS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, NZ, NA, NG, NN, NO, NZ, OM, PZ, DP, JP, JP, TP, RO, RS, RUI, SC, SD, SE, SG, SK, SI, SM, SY, TJ, TM, TN, TR, TT, TZ, DA, UG, US, UZ, VC, VW, AZ, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Burnslan (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), Buropean (AT, BE, BG, CH, CY, CZ, DE, DK, EE, BS, FL, RG, BG, RU, UI, EI, ST, LT, LU, LJ, WA, KN, NI, PL, PT, RO, SE, SI, SK, TR), OAPI (BH, BJ, CF, CG, CI, CM, GA, CM, CG), CW, MM, MM, RN, RS, NT, TD, TG).

## Published:

without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: ANTI-CTLA-4 ANTIBODY AND CPG-MOTIF-CONTAINING SYNTHETIC OLIGODEOX YNUCLEOTIDE COM-BINATION THERAPY FOR CANCER TREATMENT

(57) Abstract: The invention relates to administration of an anti-CTLA-4 antibody, perfecularly human antibodies to human CTLA-4, and the stop are the stop and t

5

10

20

25

30

Although the individual use of anti-CTLA-4 antibodies or ODNs to induce an anti-tumor response hold great promise in the treatment of cancer, there remains a need to develop novel therapies to treat tumors, more particularly, solld tumors, with such immunotherapeutic approaches.

## Summary of the Invention

Development of new therapeutic regimens, particularly those capable of augmenting or potentiating the anti-tumor activity of the immune system of the patient, while reducing the cytotoxic side effects of current chemotherapeutics, is necessary. The present invention provides such regimens.

Thus, in one embodiment, the Invention provides a method for the treatment of cancer in a patient in need of such treatment, said method comprising administering to said patient a therapeutically effective amount of an anti-CTLA-4 antibody, or antigen-binding portion thereof, in combination with a therapeutically effective amount of CpG ODN PF3512676 (CpG 7909 (also known as ProMune): TCG TCG TTT TGT CGT TTT GTC GTT; SEQ ID NO:37). In one embodiment, the 15 method is a non-vaccine method.

in one embodiment, said the CpG ODN is administered daily, every other day, twice a week, or weekly.

In one embodiment, said treatment is a therapy selected from the group consisting of necadjuvant therapy, adjuvant therapy, first-line therapy, second-line therapy, and third-line therapy.

Depending on the embodiment, said cancer is selected from the group consisting of brain cancer, breast cancer, cervical cancer, colorectal carcinoma, cutaneous T-cell lymphoma, gastric cancer, head and neck cancer, liver cancer, lung cancer, melanoma, acute myeloid leukemia. Non-Hodgkin's lymphoma, ovarian cancer, pancreatic cancer, prostate cancer, renal cell carcinoma, and sarcoma.

In other embodiments, said therapeutically effective amount of said human anti-CTLA-4 antibody ranges from about 0.1 mg/kg to 50 mg/kg, or from about 0.3 mg/kg to 20 mg/kg, including but not limited to a therapeutically effective amount of said human anti-CTLA-4 antibody selected from the group consisting of at least 1 mg/kg, at least 3 mg/kg, at least 6 mg/kg, at least 10 mg/kg, and at least 15 mg/kg.

In one embodiment, said anti-CTLA-4 antibody, or antigen-binding portion thereof, is at least one antibody selected from the group consisting of (a) a human antibody having a binding affinity for CTI A-4 of about 10-8 or greater, and which inhibits binding between CTLA-4 and B7-1, and binding between CTLA-4 and B7-2; (b) a human antibody having an amino acid sequence comprising at least one human CDR sequence that corresponds to a CDR sequence from an antibody selected from the 35 group consisting of 4.1.1, 4.8.1, 4.10.2, 4.13.1, 4.14.3, 6.1.1, 11.2.1, 11.6.1, 11.7.1., 12.3.1.1, 12.9.1.1, and 10D1; (c) a human antibody having the amino acid sequence of a heavy and/or light chain of an antibody selected from the group consisting of 4.1.1, 4.8.1, 4.10.2, 4.13.1, 4.14.3, 6.1.1, 11.2.1, 11.6.1, 11.7.1., 12.3.1.1, 12.9.1.1, and 10D1; (d) an antibody, or antigen-binding portion thereof, that competes for binding with CTLA-4 with at least one antibody having the amino acid